

## **Regional Facility Transport Protocol Summary**

As a reminder, in Southern Arizona, we have (6) regional triage protocols. You should use these, as you already do, for making ED destination decisions (regular or free-standing).

### **1. Critical Peds:**

1. Inclusion: Age < 14 y/o with altered level of consciousness, respiratory distress, or shock/hemodynamic instability
2. Destination: BUMC-TC or TMC

### **2. Trauma Patients:**

1. Inclusion: trauma and SAEMS regional Anatomic, Physiologic, or Mechanism criteria
2. Destination:
  1. Anatomic or Physiology Criteria → Level I (BUMC-TC)
  2. Mechanism criteria only or Comorbid factors → closest appropriate trauma center: BUMC-SC, BUMC-TC

### **3. Cardiac Arrest:**

1. Inclusion: Adult, cardiac arrest with return of spontaneous circulation
2. Destination: NWMC, OVMC, SMH, BUMC-TC, BUMC-SC, VA, TMC, SJH

### **4. STEMI:**

1. Inclusion: STEMI on ECG
2. Destination: NWMC, OVMC, SMH, BUMC-TC, BUMC-SC, VA, TMC, SJH

### **5. Stroke - CVA/TIA**

1. Inclusion: Cincinnati Stroke Scale (+), onset < 24hr
2. Destination: NWMC, OVMC, SMH, BUMC-TC, BUMC-SC, VA, TMC, SJH

### **6. High-Risk OB**

1. Inclusion: Preterm labor (< 37 weeks), fetal demise, premature rupture of membranes, placenta previa, placenta abruption, prolapsed cord
2. Destination: NWMC, SJH, BUMC-TC, TMC